



## Nucleotide

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search <input type="text" value="Nucleotide"/> for <input type="text"/>							<input type="button" value="Go"/>	<input type="button" value="Clear"/>
		Limits	Preview/Index	History	Clipboard	Details		
<input type="button" value="Display"/>	<input type="text" value="GenBank"/>	<input type="button" value="Save"/>	<input type="button" value="Text"/>	<input type="button" value="Add to Clipboard"/>				

1: T46645.9908 Lambda-PRL2 ...[gi:2763316]

PubMed, Taxonomy, LinkOut

LOCUS T46645 510 bp mRNA EST 04-AUG-1998  
 DEFINITION 9908 Lambda-PRL2 Arabidopsis thaliana cDNA clone 139E19T7, mRNA sequence.  
 ACCESSION T46645  
 VERSION T46645.1 GI:2763316  
 KEYWORDS EST.  
 SOURCE thale cress.  
 ORGANISM Arabidopsis thaliana  
 Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta; Spermatophyta; Magnoliophyta; eudicotyledons; core eudicots; Rosidae; eurosids II; Brassicales; Brassicaceae; Arabidopsis.  
 REFERENCE 1 (bases 1 to 510)  
 AUTHORS Newman,T., deBruijn,F.J., Green,P., Keegstra,K., Kende,H., McIntosh,L., Ohlrogge,J., Raikhel,N., Somerville,S., Thomashow,M., Retzel,E. and Somerville,C.  
 TITLE Genes galore: a summary of methods for accessing results from large-scale partial sequencing of anonymous Arabidopsis cDNA clones  
 JOURNAL Plant Physiol. 106, 1241-1255 (1994)  
 MEDLINE 95148729  
 COMMENT On Jan 9, 1998 this sequence version replaced gi:949635.  
 Contact: Thomas Newman  
 MSU-DOE Plant Research Laboratory  
 Michigan State University  
 MSU-DOE-PRL, Michigan State University, Plant Biology Bldg., E. Lansing, Mi  
 Tel: 517-353-0854  
 Fax: 517-353-9168  
 Email: 22313tcn@ibm.cl.msu.edu  
 Seq primer: T7 dye primer.  
 FEATURES  
 source  
 Location/Qualifiers  
 1..510  
 /organism="Arabidopsis thaliana"  
 /strain="var columbia"  
 /db\_xref="taxon:3702"  
 /clone="139E19T7"  
 /clone\_lib="Lambda-PRL2"  
 /note="Vector: lambda Zip-Lox; Site\_1: Sal; Site\_2: Not; Lambda PRL2 is a cDNA library derived from equal quantities of 4 pools of mRNA. The mRNA sources were 1) 7 day germinated etiolated seedlings; 2) tissue culture grown roots; 3) staged plants half with 24 hour light cycle, half on 16 hr light, 8 hour dark- rosettes; 4) same plants as 3 but aerial tissue (stems, flowers and siliques. The vector is BRL's lambda Zip-Lox. The cDNA inserts were directionally cloned with Sal-Not arms using oligo dT primed cDNA. "  
 BASE COUNT 126 a 113 c 109 g 142 t 20 others  
 ORIGIN  
 1 attttgccaa cgaacgttct ttctttcttaa tcacagcnca gcctgacgca accgctcagg  
 61 ctgatctntt ccaattttaca gccattttccc agctcagatc tctgatccgg tgagatctct  
 121 ctcaaggaaa aggagttaga gcatcatcaa gatgaaggca ctcattcttg ttggaggcct  
 181 cggaactcgc ttgagaccat tgactctcag tttcccaaag ccccttggtg atttttntaa  
 241 taaacccatg atccttcacg agatagaggc tcttaaggca gttggagttg atgaagtggg  
 301 tttggccatc aattatcagc cagaggtgat gctgaacttc ttgaaggact ttgagaccaa  
 361 gctggaaatc aaaatcactt gctcacaaga gccgagcnac taggtaccgc tggtccttgg  
 421 tttanngggn gcaaatgnt ttatggacct gngngccctt tttgttttaa aagnnatgan

481 taaggggncc cnntaaanaa ncttnggggtt  
//

Revised: October 24, 2001.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)